

**TABLE 2. Percentage distribution of pregnancies among HIV-positive female adolescents, according to selected characteristics and use of maternal health services, Kenya**

Characteristic	% (N=506)	Characteristic	% (N=506)
<b>Age</b>		<b>Unintended pregnancy</b>	
15–17	14.8	Yes	73.9
18–19	85.2	No	25.5
		Missing	0.6
<b>Study site</b>		<b>Prenatal care</b>	
Coast	14.4	Yes	84.0
Nairobi	36.2	No	15.4
Nyanza	36.8	Missing	0.6
Rift Valley	12.6		
<b>Education</b>		<b>≥4 prenatal visits</b>	
None	5.3	Yes	45.1
Primary	70.6	No	38.9
≥secondary	23.3	No visit	15.4
Missing	0.8	Missing	0.6
<b>Currently married/living together</b>		<b>PMTCT care</b>	
Yes	26.5	Yes	66.8
No	73.5	No	32.6
		Missing	0.6
<b>Age at first pregnancy</b>		<b>Pregnancy outcome</b>	
<15	11.5	Full-term live birth	74.3
15–17	67.6	Preterm live birth	10.7
18–19	20.4	Miscarriage/stillbirth	4.9
Don't remember/missing	0.6	Ended it/aborted	4.2
		Still pregnant	5.3
<b>Pregnancy/order</b>		Missing	0.6
1	73.7	<b>Skilled attendance‡</b>	
2	21.7	Yes	72.9
3	3.6	No	27.1
4	1.0		
<b>Man involved in pregnancy</b>		<b>Postnatal/postabortion care‡</b>	
Boyfriend/fiancé	66.2	Yes	82.8
Husband	24.7	No	17.2
Other†	8.7		
Missing	0.4	Total	100.0

†Friend, acquaintance or stranger. ‡Among the 476 pregnancies that had ended (live births, stillbirths, miscarriages or abortions). Notes: PMTCT=prevention of mother-to-child transmission (of HIV). Percentages may not total 100.0 because of rounding.