

TABLE 3. Selected sexual behaviors in various subsamples of partnerships, by gender

| Measure | All partnerships | | Restricted partnerships | | All partnerships (adjusted) | | Matched couples | |
|--------------------------------------|------------------|-------------|-------------------------|-------------|-----------------------------|-------------|-----------------|-------------|
| | Women (N=549) | Men (N=750) | Women (N=157) | Men (N=330) | Women (N=549) | Men (N=750) | Women (N=280) | Men (N=280) |
| Time until first sex (mos.) | 4.9 | 3.7* | 3.8 | 3.2 | 4.8 | 3.7 | 3.4 | 3.4 |
| Frequency of sex (first month) | | | | | | | | |
| 1–4 times | 57.5 | 62.9* | 57.4 | 62.6 | 59.0 | 61.5 | 10.1 | 10.8 |
| 5–14 times | 30.8 | 29.8 | 34.8 | 29.2 | 31.2 | 29.7 | 22.7 | 30.6 |
| ≥15 | 11.8 | 7.3 | 7.7 | 8.2 | 9.7 | 8.9 | 67.2 | 58.6 |
| Frequency of sex (last month) | | | | | | | | |
| 0 | 31.4 | 41.6*** | 24.4 | 31.7 | 35.7 | 38.2 | 16.9 | 17.9 |
| 1–4 times | 31.8 | 34.0 | 34.0 | 37.2 | 33.2 | 32.9 | 27.0 | 23.3 |
| 5–14 times | 24.5 | 16.3 | 29.5 | 20.1 | 20.6 | 19.4 | 39.6 | 40.5 |
| ≥15 | 12.4 | 8.2 | 12.2 | 11.0 | 10.5 | 9.6 | 16.6 | 18.3 |
| Used any contraceptive (first month) | 67.0 | 71.4 | 69.4 | 72.5 | 72.0 | 67.4 | 62.8 | 62.5 |
| Used condoms (first month) | | | | | | | | |
| Rarely/never | 43.7 | 31.4*** | 41.0 | 31.6** | 38.7 | 34.9 | 43.7 | 46.4 |
| Sometimes/mostly | 15.7 | 13.0 | 18.0 | 11.9 | 14.5 | 14.1 | 16.9 | 16.4 |
| Always | 40.6 | 55.6 | 41.0 | 56.5 | 46.9 | 51.0 | 39.4 | 37.1 |
| Used condoms (last month) | | | | | | | | |
| No sex | 31.6 | 41.1*** | 25.8 | 31.4*** | 36.2 | 37.5 | 17.6 | 20.4 |
| Rarely/never | 43.5 | 23.7 | 47.7 | 28.1 | 32.0 | 31.9 | 49.6 | 50.4 |
| Sometimes/mostly | 8.2 | 6.6 | 9.7 | 7.0 | 7.4 | 7.3 | 11.5 | 11.1 |
| Always | 16.6 | 28.7 | 16.8 | 33.5 | 24.3 | 23.4 | 21.2 | 18.2 |
| Ever used condoms | 64.5 | 75.7*** | 68.0 | 76.4* | 70.1 | 71.7 | 62.7 | 60.7 |
| Man had other nonmarital partner† | 17.3 | 25.1*** | 17.2 | 20.9 | 20.4 | 22.4 | 10.7 | 12.9 |
| Woman had other nonmarital partner‡ | 6.7 | 25.1*** | 6.4 | 25.5*** | 9.5 | 21.0*** | 3.9 | 8.2* |

* $p < .05$. ** $p < .01$. *** $p < .001$. †Compares men's reports of infidelity with women's suspicion of their partner's infidelity. ‡Compares women's reports of infidelity with men's suspicion of their partner's infidelity. Notes: "First month" refers to the first month of sexual activity in the relationship. Chi-square and t tests were used to assess differences in the all-partnerships and restricted-partnerships samples; Z tests were used for the all-partnerships adjusted sample; and McNemar, Wilcoxon signed rank and chi-square tests were used for the matched-couples sample.