

TABLE 2. Percentage of medication abortion clients for whom assessments of eligibility and abortion completion status differed between study provider and verifier, and related measures—all by provider type; and differences between provider types in accuracy of assessments (and 95% confidence intervals)

Indicator	Provider type			Difference	
	Ayurvedic physicians	Nurses	Allopathic physicians	Ayurvedic vs. allopathic	Nurses vs. allopathic
ELIGIBILITY					
No. screened	461	497	456	na	na
Provider's assessment differed from verifier's	3.3	4.0	3.9	-0.7 (-3.1 to 1.7)	0.1 (-2.4 to 2.6)
Provider classified woman as ineligible, verifier as eligible	1.3	2.0	0.7	0.6 (-0.6 to 1.9)	1.4 (-0.1 to 2.8)
Provider classified woman as eligible, verifier as ineligible	2.0	2.0	3.3	-1.3 (-3.4 to 0.7)	-1.3 (-3.3 to 0.8)
Sensitivity	98.5	97.6	99.3	na	na
Specificity	84.2	87.7	70.6	na	na
ABORTION COMPLETENESS					
No. of assessments	412	433	435	na	na
Provider's assessment differed from verifier's	3.6	5.1	4.4	-0.7 (-3.4 to 1.9)	0.7 (-2.1 to 3.5)
Provider considered abortion incomplete, verifier complete	1.5	3.0	2.1	-0.6 (-2.4 to 1.2)	0.9 (-1.2 to 3.0)
Provider considered abortion complete, verifier incomplete	2.2	2.1	2.3	-0.1 (-2.1 to 1.9)	-0.2 (-2.2 to 1.7)
Sensitivity	98.3	96.5	97.6	na	na
Specificity	82.4	84.2	84.1	na	na

Notes: Equivalence is indicated when the 95% confidence interval for the difference in failure rates between the allopathic physician group and each of the other two groups (ayurvedic physicians and nurses) lies within the predetermined margin of equivalence ($\pm 5.5\%$). na=not applicable.