

**TABLE 3. Odds ratios (and 95% confidence intervals) from logistic regression analyses examining the likelihood that sexually active pill users missed more than one pill and used no contraceptive backup for intercourse subsequent to the last missed pill during the four weeks prior to interview, by selected characteristics**

Characteristic	Missed > 1 pill and used no backup (N=54)
<b>Age</b>	
18–24	0.9 (0.3–3.1)
25–34	1.0
35–44	0.5 (0.2–1.3)
<b>Level of education</b>	
<H.S.	1.0
≥H.S.	1.0 (0.4–2.7)
<b>Unemployed</b>	
No	1.0
Yes	1.2 (0.5–3.2)
<b>Nationality</b>	
French	1.0
Other	2.6 (0.9–7.3)†
<b>Age of the last child</b>	
No children	1.0
<2 years	3.0 (1.0–9.3)
≥2 years	1.9 (0.7–5.4)
<b>History of unintended pregnancy</b>	
Yes	2.7 (1.2–6.1)*
No	1.0
<b>Partner at last intercourse</b>	
Regular	1.0
Occasional	4.8 (0.5–49.1)
<b>Involved in the choice of the pill</b>	
Yes	1.0***
No	11.7 (3.2–42.4)
<b>Established pill-taking routine</b>	
Yes	1.0**
No	4.3 (1.7–10.6)
<b>Health side effects</b>	
Yes	1.5 (0.6–4.1)
No	1.0
<b>Negative effect of the pill on sexual desire</b>	
Yes	0.8 (0.2–3.1)
No	1.0
<b>Type of pill</b>	
Combined	1.0*
Progestin-only	2.5 (1.0–6.2)
<b>Duration of use</b>	
≤1 year	0.8 (0.3–2.1)
>1 year	1.0

\*p<.05. \*\*p<.01. \*\*\*p<.001. †p<.10. Note: Significance levels refer to the results for the likelihood ratio test for all the categories of a variable.