

**TABLE 4. Percentage of women with correct knowledge of contraceptive methods and healthy pregnancy spacing messages, at baseline, four months postpartum and nine months postpartum, by study group**

Topic	Intervention (N=477)			Comparison (N=482)		
	Baseline	4 mos. postpartum	9 mos. postpartum	Baseline	4 mos. postpartum	9 mos. postpartum
Knew IUD is placed in uterus	44.2	66.0**	84.3**	30.6	31.2	45.8
Knew IUD is effective for 10 yrs.	0.4	29.9**	47.8**	0.4	2.5	5.4
Knew correct pill use	60.6	79.5	92.2*	53.9	75.6	86.3
Knew correct condom use	81.6	93.0*	94.3*	80.9	84.0	85.5
Knew correct emergency contraceptive use	0.9	8.6**	19.7**	1.2	0.2	3.7
Knew all 3 conditions required for lactational amenorrhea method	0.0	64.4**	na	0.0	0.2	na
Mentioned 3–5 yrs. spacing between births	86.6	92.2	na	89.8	91.6	na
Mentioned at least 3 undesirable outcomes of short-spaced pregnancy	32.1*	55.4**	na	38.2	25.5	na
Mentioned health of woman, last child and fetus affected by short-spaced pregnancy	30.2	70.0*	na	28.4	63.7	na

\*p≤.05. \*\*p≤.01. Note: na=not applicable.