

**TABLE 7. Unadjusted and adjusted odds ratios from logistic regression analyses identifying associations between selected characteristics and knowledge of healthy spacing practices at four months postpartum and of spacing methods at nine months postpartum**

Characteristic	Knowledge of healthy spacing messages		Knowledge of $\geq 2$ spacing methods	
	Unadjusted odds ratio	Adjusted odds ratio	Unadjusted odds ratio	Adjusted odds ratio
<b>Study group</b>				
Comparison (ref)	1.00	1.00	1.00	1.00
Intervention	2.18**	2.07**	2.06**	1.96**
<b>Baseline value</b>	0.98	0.97	2.00**	1.53*
<b>Woman's education</b>				
Illiterate (ref)	na	1.00	na	1.00
Primary school	na	0.96	na	1.11
Middle school	na	1.11	na	2.07**
$\geq$ high school	na	1.31	na	3.18**
<b>Age</b>				
15–19 (ref)	na	1.00	na	1.00
20–24	na	1.65*	na	1.46
<b>Age at initiation of cohabitation</b>				
15–19 (ref)	na	1.00	na	1.00
20–23	na	0.93	na	0.77
<b>Caste</b>				
Scheduled caste/tribe (ref)	na	1.00	na	1.00
Other backward-caste Hindu	na	1.27	na	1.26
High-caste Hindu	na	1.93**	na	1.66
Other religious groups	na	1.41	na	1.39
<b>Parity</b>				
0 (ref)	na	1.00	na	1.00
1	na	0.89	na	0.82
<i>-log likelihood</i>		490.25		382.59
<i>Pseudo R<sup>2</sup>/R<sup>2</sup></i>		0.04		0.08

\* $p \leq .05$ . \*\* $p \leq .01$ . Notes: ref=reference group. na=not applicable.