

TABLE 3. Selected characteristics of Asian and Pacific Islander young adults, by reported STD history

Characteristic	Ever had an STD	Never had an STD	Characteristic	Ever had an STD	Never had an STD
PERCENTAGE DISTRIBUTIONS			Condom used at last sex		
Gender			Yes	82.4	75.7
Male	24.7	51.7	No	17.4	24.3
Female	75.3	48.4	<i>F(df)=1.09 (1,122)</i>		
<i>F(df)=10.50 (1,123)***</i>			Ever paid for sex		
Ethnicity			Yes	4.9	2.5
Chinese	20.5	19.9	No	95.1	97.6
Filipino	14.1	22.9	<i>F(df)=2.7 (1,122)†</i>		
Japanese	5.3	8.6	Ever been paid for sex		
Indian	28.2	11.2	Yes	8.6	1.8
Korean	13.6	15.3	No	91.4	98.2
Vietnamese	13.3	20.8	<i>F(df)=12.8 (1,122)***</i>		
Other	9.8	9.6	Had sex before age 15		
<i>F(df)=2.10 (1,123)†</i>			Yes	30.5	14.6
Age			No	69.5	85.4
18	0.0	0.1	<i>F(df)=534 (1,123)*</i>		
19	6.6	8.3	Had >1 sex partner in past 12 mos.		
20	28.1	13.0	Yes	55.4	27.1
21	15.3	14.5	No	44.6	72.9
22	20.7	17.5	<i>F(df)=12.1 (1,121)***</i>		
23	12.8	19.9	Total	100.0	100.0
24	14.0	18.5	MEANS		
25	7.1	2.5	Health Belief Model		
26	0.0	8.2	Perceived susceptibility	1.51 (0.94)	1.17 (0.54)
27	0.0	3.5	<i>t(df)=5.39 (1,094)***</i>		
<i>F(df)=1.08 (1,123)</i>			Perception of responsiveness to treatment	4.47 (0.97)	4.10 (1.12)
Acculturation			<i>t(df)=3.19 (1,094)***</i>		
Speak English at home, U.S.-born	56.6	48.5	Perception of relationship consequences	4.26 (1.26)	4.54 (0.99)
Speak English at home, foreign-born	39.8	41.2	<i>t(df)=2.57 (1,048)**</i>		
Speak no English at home, U.S.-born	3.1	5.4	External cues to action	4.20 (1.08)	3.96 (1.11)
Speak no English at home, foreign-born	0.5	4.9	<i>t(df)=2.10 (1,074)*</i>		
<i>F(df)=1.14 (1,123)</i>			Internal cues to action	4.27 (1.06)	4.16 (1.05)
Binge drinking in past 12 mos.			<i>t(df)=1.00 (1,086)</i>		
Yes	76.0	70.0			
No	24.0	30.0			
<i>F(df)=0.94 (1,119)</i>					

*p<.05. **p<.01. ***p<.001. †p<.10. Notes: Percentages incorporate population-based sampling weights. Because of imprecision in the estimated weights, some variables may contain weighted percentages that do not add to 100%. For the survey adjusted chi-square test, p values are calculated from an approximate F distribution. Figures in parentheses accompanying means are standard deviations.