

TABLE 4. Odds ratios from logistic regression analyses identifying associations of HIV status and women's characteristics with use of contraceptives at time of unintended pregnancy in the last five years, pooled sample

| Characteristic | Model 1 | Model 2 |
|---|---------|---------|
| HIV status | | |
| Known, positive | 2.84† | 3.24* |
| Known, negative (ref) | 1.00 | 1.00 |
| Unknown | 0.50* | 0.46* |
| Country | | |
| Nigeria | 0.54 | 0.49 |
| Zambia (ref) | 1.00 | 1.00 |
| Interaction terms | | |
| Nigeria x HIV-positive | 0.93 | 0.40 |
| Nigeria x unknown HIV status | 1.34 | 1.76 |
| Age | | |
| 18–24 (ref) | na | 1.00 |
| 25–34 | na | 2.37* |
| 35–49 | na | 0.95 |
| Residence | | |
| Urban | na | 0.58* |
| Rural (ref) | na | 1.00 |
| Highest level of school attended | | |
| None | na | 0.48 |
| Primary | na | 0.78 |
| ≥secondary (ref) | na | 1.00 |
| Union status | | |
| Married or cohabiting | na | 1.32 |
| Not in union (ref) | na | 1.00 |
| No. of living children | | |
| 0 | na | 1.15 |
| 1–2 | na | 0.68 |
| 3–4 | na | 0.71 |
| ≥5 (ref) | na | 1.00 |

* $p \leq .05$. † $p \leq .10$. Notes: Both models were estimated for the 327 women who had had an unintended pregnancy in the last five years. ref=reference group. na=not applicable. Ns are unweighted.