

TABLE 5. Odds ratios from logistic regression analyses identifying associations of HIV status and women's characteristics with having had an abortion in the last five years, pooled sample

| Variable | Model 1 | Model 2 |
|---|---------|---------|
| HIV status | | |
| Known, positive | 2.82 | 2.04 |
| Known, negative (ref) | 1.00 | 1.00 |
| Unknown | 4.02* | 3.76† |
| Country | | |
| Nigeria | 8.23** | 4.42† |
| Zambia (ref) | 1.00 | 1.00 |
| Interaction terms | | |
| Nigeria x HIV-positive | 1.70 | 2.17 |
| Nigeria x unknown HIV status | 0.70 | 0.62 |
| Age | | |
| 18–24 (ref) | na | 1.00 |
| 25–34 | na | 1.04 |
| 35–49 | na | 1.57 |
| Residence | | |
| Urban | na | 2.89* |
| Rural (ref) | na | 1.00 |
| Highest level of school attended | | |
| None | na | 0.77 |
| Primary | na | 0.75 |
| ≥secondary (ref) | na | 1.00 |
| Union status | | |
| Married or cohabiting | na | 0.82 |
| Not in union (ref) | na | 1.00 |
| No. of living children | | |
| 0 | na | 11.35** |
| 1–2 | na | 1.91 |
| 3–4 | na | 2.31 |
| ≥5 (ref) | na | 1.00 |

* $p \leq .05$. ** $p \leq .01$. † $p \leq .10$. Notes: Both models were estimated for the 327 women who had had an unintended pregnancy in the last five years. ref=reference group. na=not applicable. Ns are unweighted.