

**TABLE 3. Percentage distribution of women using reversible contraceptives, by current method used, according to pregnancy-, method- and provider-related experiences and attitudes; and odds ratios from logistic regression analyses examining associations between experiences and attitudes and use of specific methods**

Experience or attitude	N	%					Odds ratio			
		Pill	Long-acting	Condom	Other	Total	Pill	Long-acting	Condom	Other
ALL	1,641	37.7	17.9	32.3	12.1	100.0	na	na	na	na
<b>PREGNANCY</b>										
<b>Parity</b>										
0 (ref)	637	49.8	11.4	31.2	7.5	100.0	1.00	1.00	1.00	1.00
1	377	32.1†	21.8†	31.6	14.6†	100.0	0.81	1.67*	0.87	1.20
≥2	626	28.8†	21.9†	33.8	15.5†	100.0	0.70	1.97**	0.87	1.19
<b>No. of unintended pregnancies</b>										
0 (ref)	976	43.9	12.6	33.2	10.3	100.0	1.00	1.00	1.00	1.00
≥1	664	28.8†	25.6†	30.9	14.8†	100.0	0.73*	2.10***	0.66**	1.30
<b>Importance of avoiding pregnancy</b>										
Very (ref)	1,045	40.5	18.8	33.7	7.1	100.0	1.00	1.00	1.00	1.00
Somewhat	335	41.5	17.3	25.7†	15.5†	100.0	0.96	0.91	0.83	2.01***
A little/not	261	21.9†,‡	15.0	35.0	28.1†,‡	100.0	0.38***	0.77	1.14	4.42***
<b>Fatalistic attitude toward pregnancy/birth control</b>										
Disagree/neutral (ref)	1,168	40.9	17.7	31.9	9.5	100.0	1.00	1.00	1.00	1.00
Agree	473	30.0†	18.2	33.2	18.6†	100.0	0.96	0.82	0.82	1.91***
<b>METHOD AND PROVIDER</b>										
<b>Reason for method use</b>										
Mostly like method/both like and dislike (ref)	1,017	46.1	21.9	21.9	10.0	100.0	1.00	1.00	1.00	1.00
Mostly don't like other methods	624	24.1†	11.2†	49.1†	15.6†	100.0	0.37***	0.42***	4.04***	1.42*
<b>Would change method if cost were not an issue</b>										
No (ref)	1,134	42.6	19.6	27.0	10.7	100.0	1.00	1.00	1.00	1.00
Yes	506	26.8†	13.8†	44.0†	15.4†	100.0	0.58***	0.46***	2.18***	1.90***
<b>Type of provider</b>										
Private doctor (ref)	1,031	41.8	16.8	28.8	12.6	100.0	1.00	1.00	1.00	1.00
Clinic	472	38.0	23.1†	29.3	9.6	100.0	1.06	1.05	0.90	0.97
None	137	6.6†,‡	7.3†,‡	68.6†,‡	17.5‡	100.0	0.10***	0.33**	4.71***	1.57
<i>Model R<sup>2</sup> (Nagelkerke)</i>	<i>na</i>	<i>na</i>	<i>na</i>	<i>na</i>	<i>na</i>	<i>na</i>	<i>0.269</i>	<i>0.194</i>	<i>0.267</i>	<i>0.217</i>

\*p<.05. \*\*p<.01. \*\*\*p<.001. †Significantly different from percentage in first row at p<.05. ‡Significantly different from percentage in second row at p<.05. Notes: Ns are weighted. Regressions include all variables listed in Table 2. Long-acting methods are the injectable, patch, IUD, ring and implant. Other methods are withdrawal, periodic abstinence, spermicides and other barrier methods. na=not applicable. ref=reference group.