

**TABLE 5. Percentage of women using oral contraceptives or condoms who reported inconsistent use, by pregnancy-, method- and provider-related experiences and attitudes; and odds ratios from logistic regression analyses examining associations between experiences and attitudes and inconsistent method use**

Experience or attitude	Pill		Condom		Odds ratio	
	N	%	N	%	Pill	Condom
ALL	619	38.2	529	61.1	na	na
<b>PREGNANCY</b>						
<b>Parity</b>						
0 (ref)	318	42.6	199	58.8	1.00	1.00
1	121	41.3	119	65.5	1.06	1.04
≥2	180	28.3†	211	60.7	0.50*	0.82
<b>No. of unintended pregnancies</b>						
0 (ref)	429	38.0	205	63.4	1.00	1.00
≥1	191	38.4	324	57.6	1.40	0.57*
<b>Importance of avoiding pregnancy</b>						
Very (ref)	423	39.0	352	57.7	1.00	1.00
Somewhat	139	38.1	86	58.6	1.16	0.92
A little/not	57	32.8	91	76.7†,‡	0.86	2.58**
<b>Fatalistic attitude toward pregnancy/birth control</b>						
Disagree/neutral (ref)	477	40.0	372	57.3	1.00	1.00
Agree	142	32.4	157	70.1†	0.81	1.48
<b>METHOD</b>						
<b>Dual method use</b>						
No (ref)	431	35.7	384	49.0	1.00	1.00
≥2 methods together	130	40.5	51	82.4†	1.26	6.33***
≥2 methods switching	58	50.8	94	98.9†,‡	1.45	77.79***
<b>Duration of current method use (in yrs.)</b>						
<2	144	48.6	163	68.7	1.78*	1.91*
2–4	175	34.9†	130	59.7	0.89	1.42
≥5 (ref)	296	34.8†	232	56.0†	1.00	1.00
<b>Satisfied with method in past year</b>						
Completely (ref)	456	34.9	242	55.4	1.00	1.00
Not completely	161	48.1†	286	66.1†	1.56*	1.91**
<b>PROVIDER</b>						
<b>Type of provider</b>						
Private doctor (ref)	431	37.4	297	57.7	1.00	1.00
Clinic	179	40.2	138	67.6	1.01	1.39
None	9	44.4	94	61.7	0.43	1.43
<b>Satisfied with provider</b>						
Very (ref)	403	33.5	326	61.0	1.00	1.00
Not very	216	46.8†	203	61.1	1.59*	0.94
<b>Usually see same clinician</b>						
Yes/ no visit (ref)	514	35.6	431	60.3	1.00	1.00
No	106	50.9†	98	64.6	1.67*	0.99
<i>Model R<sup>2</sup> (Nagelkerke)</i>	<i>na</i>	<i>na</i>	<i>na</i>	<i>na</i>	<i>0.159</i>	<i>0.405</i>

\*p<.05. \*\*p<.01. \*\*\*p<.001. †Significantly different from percentage in first row at p<.05. ‡Significantly different from percentage in second row at p<.05. Notes: Women who reported having used more than one method were classified according to their most effective method. Inconsistent pill use is defined as missing one or more pills in prior three months; inconsistent condom use is defined as not using a condom at each sex in prior three months or putting the condom on late at least once. Ns are weighted. Regressions include all variables listed in Table 4. na=not applicable. ref=reference group.