

TABLE 2. Odds ratios from logistic and multinomial logistic regression analyses examining relationship of time period and interaction between time period and area with pregnancy termination and pregnancy termination method

| Measure | Model 1‡ | Model 2§ | | Model 3‡ |
|----------------------------------|------------------------------|---|---|--|
| | Whether pregnancy terminated | Whether pregnancy terminated by safer method (relative to not terminated) | Whether pregnancy terminated by less-safe method (relative to not terminated) | Whether termination was by safer method (relative to less-safe method) |
| Time period | | | | |
| 1989–1993 (ref) | 1.00 | 1.00 | 1.00 | 1.00 |
| 1994–1998 | 1.25*** | 1.63*** | 0.96 | 1.68*** |
| 1999–2003†† | 1.79*** | 2.96*** | 0.85† | 3.56*** |
| 2004–2008 | 1.35*** | 2.35*** | 0.52* | 4.54*** |
| Time period x MCH-FP area | | | | |
| 1989–1993 x MCH-FP area | 0.66*** | 0.67*** | 0.66*** | 0.97 |
| 1994–1998 x MCH-FP area | 0.41*** | 0.48*** | 0.32*** | 1.49** |
| 1999–2003 x MCH-FP area†† | 0.47*** | 0.46*** | 0.48*** | 0.91 |
| 2004–2008 x MCH-FP area | 0.62*** | 0.63*** | 0.57*** | 1.07 |

*p<.05. **p<.01. ***p<.001. †p<.10. ‡Logistic regression. §Multinomial logistic regression. ††Data not available for 2001. Notes: Odds ratios for maternal age and education are same as in Table 1. Models 1 and 2 include all eligible pregnancies; model 3 is restricted to pregnancies that were terminated. ref=reference category. MCH-FP=Maternal and Child Health–Family Planning.