

TABLE 5. Odds ratios (and 95% confidence intervals) from multinomial logistic regression analyses identifying associations between respondent characteristics and the likelihood of having used emergency contraceptive pills 2–5 or six or more times over the last six months, rather than no more than once, Lagos

Characteristic	Unadjusted		Adjusted	
	2–5 times (N=173)	≥6 times (N=142)	2–5 times (N=134)	≥6 times (N=112)
Age†				
18–34 (ref)	1.00	1.00	1.00	1.00
35–49	0.42 (0.26–0.70)**	0.36 (0.23–0.57)**	0.68 (0.30–1.55)	0.53 (0.23–1.25)
Marital status				
Single (ref)	1.00	1.00	1.00	1.00
Ever-married†	0.62 (0.42–0.94)*	0.68 (0.39–1.20)	0.59 (0.29–1.22)	0.86 (0.37–2.01)
Educational attainment				
≤secondary† (ref)	1.00	1.00	1.00	1.00
≥college	1.14 (0.78–1.67)	1.54 (0.75–3.15)	0.95 (0.44–2.08)	1.52 (0.53–4.32)
Socioeconomic status				
Low (ref)	1.00	1.00	1.00	1.00
Middle	0.82 (0.50–1.36)	0.91 (0.48–1.74)	0.58 (0.24–1.38)	0.41 (0.14–1.22)
High	1.46 (0.79–2.68)	0.62 (0.27–1.42)	0.75 (0.39–1.42)	0.28 (0.14–0.56)**
Frequency of intercourse				
<once a week (ref)	1.00	1.00	1.00	1.00
Once a week	5.63 (1.81–17.52)**	13.75 (3.19–59.31)**	4.38 (1.49–12.85)**	10.10 (2.01–50.77)**
2–3 times a week	10.23 (2.53–41.29)**	11.14 (2.18–56.94)**	10.29 (2.49–42.48)**	14.32 (2.10–97.67)**
≥4 times a week	7.50 (0.95–59.46)	25.00 (5.25–118.95)**	7.76 (1.18–50.95)*	31.58 (5.56–179.41)**
Used condoms as main method				
No (ref)	1.00	1.00	1.00	1.00
Yes	1.48 (0.83–2.65)	0.93 (0.45–1.92)	0.63 (0.27–1.43)	0.37 (0.18–0.78)**
Used the pill/injectable as main method				
No (ref)	1.00	1.00	1.00	1.00
Yes	0.60 (0.31–1.18)	0.28 (0.12–0.64)**	0.37 (0.14–0.96)*	0.12 (0.04–0.37)**
Used natural method as main method				
No (ref)	1.00	1.00	1.00	1.00
Yes	0.53 (0.28–1.02)	0.27 (0.10–0.74)*	0.27 (0.09–0.78)*	0.15 (0.04–0.62)**

*p<.05. **p<.01. †Because of small cell sizes, several categories were combined. Notes: In the unadjusted and adjusted models, the comparison groups of women who had used the pills no more than once in the last six months included 128 and 70 individuals, respectively. The overall N in the adjusted model was greatly reduced because of missing observations for the independent variables. ref=reference group.