Table 4. Among married women currently using a method to space births, estimated average duration of contraceptive use and percentage who terminate use each year, by country

Country	Months of use	% terminating
Peru	64.2	18.7
Indonesia	62 6	19 2

•			•
Peru	64.2	18.7	
Indonesia	62.6	19.2	
Colombia	57.2	21.0	
Paraguay	57.2	21.0	

Peru	04.2	10.7	
Indonesia	62.6	19.2	
Colombia	57.2	21.0	
Paraguay	57.2	21.0	
Bolivia	55.2	21.8	
Turkey	48.4	24.8	
7' ' '	4-0		

Colombia	57.2	21.0	
Paraguay	57.2	21.0	
Bolivia	55.2	21.8	
Turkey	48.4	24.8	
Zimbabwe	47.6	25.2	
Philippines	46.2	26.0	

Bolivia	55.2	21.8	
Turkey	48.4	24.8	
Zimbabwe	47.6	25.2	
Philippines	46.2	26.0	
Morocco	41.1	28.9	
Eavpt	40.0	30.0	

Zimbabwe	47.6	25.2	
Philippines	46.2	26.0	
Morocco	41.1	28.9	
Egypt	40.0	30.0	
Dominican Pon	33 U	26.5	

ominican Rep. 33.U 36.5

Note: The estimated duration of use is twice the current duration, assuming that women are surveyed, on average, halfway through an in-

terval of use. The termination rate is the reciprocal of total years of use.