

Table 3. Linear logistic regression coefficients and adjusted odds ratios (and 95% confidence intervals) indicating the likelihood of nonuse of contraceptives

Variable	Coefficient	Standard error	p	Adjusted odds ratio
Place of last delivery				
Home	0.8309	0.1611	.0001	2.30 (1.67–3.15)
Medical facility	na	na	na	1.00
Age at first pregnancy				
≤16	na	na	na	1.00
17–19	0.5422	0.1945	.0053	1.72 (1.18–2.52)
≥20	1.0134	0.2175	.0001	2.76 (1.80–4.22)
No. of children who died				
0	na	na	na	1.00
1	0.1453	0.2280	.5239	1.16 (0.74–1.81)
≥2	1.0367	0.2743	.0002	2.82 (1.65–4.83)
Family structure				
Extended	0.8979	0.1987	.0001	2.45 (1.66–3.62)
Nuclear	na	na	na	1.00
Currently employed				
Yes	na	na	na	1.00
No	0.7831	0.2581	.0024	2.19 (1.32–3.63)
Type of floors				
Mud	0.4874	0.1601	.0023	1.63 (1.19–2.23)
Covered	na	na	na	1.00
Education				
None	0.4371	0.1930	.0236	1.55 (1.06–2.26)
Some primary	–0.2601	0.2755	.3451	0.77 (0.45–1.32)
Complete primary	–0.1525	0.2869	.5951	0.86 (0.49–1.51)
≥secondary	na	na	na	1.00
<i>Intercept</i>	<i>–2.8619</i>	<i>0.3249</i>	<i>.0001</i>	<i>na</i>

Notes: All variables have a .05 significance level for entry into the model, except two education variables and one child mortality variable. Data are missing for 15 women (N=868). na=not applicable.