

Table 3. Odds ratios (and 95% confidence intervals) showing the likelihood that clients used an effective contraceptive at last intercourse, by clients' characteristics

| Characteristic | Odds ratio |
|----------------------------------|-------------------|
| Demographic | |
| White vs. Hispanic | 1.10 (0.85–1.40) |
| White vs. black | 1.11 (0.80–1.54) |
| White vs. Asian | 1.07 (0.74–1.57) |
| Aged 18–19 vs. <14 | 0.97 (0.67–1.40) |
| Aged 18–19 vs. 15–17 | 0.96 (0.78–1.22) |
| Receives public assistance† | 0.71 (0.50–1.02) |
| Sexual behavior | |
| Not comfortable with method | 0.40* (0.33–0.50) |
| Agree with partner about method | 1.39* (1.06–1.81) |
| Never got someone pregnant | 1.89* (1.46–2.43) |
| Never had STD | 0.85 (0.58–1.24) |
| Never had HIV test | 0.88 (0.65–1.26) |
| Psychosocial | |
| No problems in school | 1.07 (0.84–1.40) |
| No signs of depression | 1.09 (0.79–1.52) |
| High during sex | 0.96 (0.75–1.21) |
| Problems talking with others | 0.88 (0.59–1.31) |
| Unemployed | 1.09 (0.83–1.42) |
| Problems with gangs | 1.35 (0.77–2.36) |
| Not homeless or runaway | 0.59 (0.28–1.24) |
| Never forced or tricked into sex | 0.78 (0.49–1.29) |
| No teenage parent in family | 1.08 (0.86–1.36) |

* $p < .05$. †Medicaid or Aid to Families with Dependent Children.