

Table 4. Percentage distribution of providers, by proportion of patients with HPV to whom they offer counseling and educational interventions, according to intervention (N=140)

Intervention	% of patients					Total
	100	75	50	1–50	0	
Take a sexual history on new HPV patients	77.1	13.6	4.3	3.6	1.4	100.0
Inquire about smoking status	59.3	15.0	10.7	7.1	7.9	100.0
Discuss potential of HPV recurrence*	89.4	8.5	0.7	1.4	0.0	100.0
Discuss potential for malignant changes	85.2	12.7	0.7	1.4	0.0	100.0
Instruct on genital self-exams*,†	53.9	24.8	12.8	5.0	3.5	100.0
Discuss issues of body image/self-esteem*,‡	40.4	31.9	14.9	9.2	3.5	100.0
Discuss feelings of anger, guilt and blame*,‡,§	47.9	33.1	8.5	7.0	3.5	100.0
Give out printed materials*,†,‡	71.8	19.7	2.8	4.9	0.7	100.0
Role-play‡	5.0	5.7	10.6	25.5	53.2	100.0
Refer to support group	1.4	6.4	12.8	24.1	55.3	100.0
Do joint counseling with partner	2.1	4.3	14.9	53.9	24.8	100.0

*Nurse practitioners reported the provision of this intervention to 100% of their patients with significantly greater frequency than physicians ($p < .05$). †Nurse practitioners reported the provision of this intervention to 100% of their patients with significantly greater frequency than female physicians ($p < .005$). ‡Female providers reported the provision of this intervention to 100% of their patients with significantly greater frequency than male providers ($p < .05$). §Female physicians reported the provision of this intervention to 100% of their patients with significantly greater frequency than male physicians ($p < .05$).