Characteristic Female Male Interven-

who were surveyed at pretest, posttest and one-year follow-up, by sex

	(N=103)	(N=107)		(N=22)	
Mean age	12.8	12.9	.817	13.1	
Mean grade	7.3	7.4	.301	7.3	
% black/Caribbean	70.3	62.7	.267	64.7	
% Hispanic/Latino	19.8	23.5	.531	17.6	

18.6

45.3

245

25.4

16.8

18.4

214

87

18.0

10.8

0.0

63.0

44.6

*The mean depression score is based on a scale of six items, with possible scores ranging from 6 to 18.

3.0

% multiethnic/other

in last month

% who repeated a grade

Mean depression score*

% who cut class at least once

% who live in two-parent household

% who report having been slapped, punched or kicked by a parent/quardian

% who used alcohol in past 30 days

% who have been touched sexually when it was not desired

% who have been forced to have sex

% who were sexually active at pretest

% who have ever been pregnant or have ever caused a pregnancy

% who report that at least one friend who has been pregnant or has ever caused a pregnancy

% who report that a few/most/all of their friends have had sex

% who report they will definitely/probably have sex in the next 6 months

% who used cigarettes in past 30 days

% who used marijuana in past 30 days

% who live in a household where English is spoken about half of the time or less

tion

19.8 .531

Compar-

ison

Table 3. Selected means and percentages for characteristics at pretest of study participants

9.9 13.8 .696 18.4 7.5 .020

15.8 11.27 10.24

39 4

25.2

9.9

11.5

11.5

47

6.5

28

19.8

8.5

0.0

58 1

43.3

.602 .009 257

429

.017

.269

.177

545

002

064

.742

.577

1 000

462

.841

р

(N=22)	
13.1 7.3 64.7 17.6 17.6 27.3	
27.2 10.33 44.4	

59

12.5

10.0

5.0

95

0.0

0.0

44.5

31.8

0.0

65.0

33.4

Interven-

tion

Compar-

ison

(N=80)12.9

74 68.0

26.7

13.9

15.2

50.0

25.0

17.0

11.4

5.1

64

38

1.3

42.7

20.1

13

49 4

32.9

10.25

5.3

р

.292 852

796

444

.181

.213

.194

.996

539

017

.673

.982

.861

625

358

.600

.893

.296

384

216

.970