care visits for their last live birth in the previous five years, by dimensions of deprivation; and ratio of percentages, by dimensions of deprivation—according to state India/state ΑII Dimensions of deprivation Ratio of none to

Two

51.7

46.9

50.0

26.0

91.2

14.8

30.2

48.5

Three

29.7

47.1

15.1

u

u

8.9

u

44.7

One

1.1

1.0

1.3

1.2

1.1

1.3

1.4

1.1

1.7

1.8

1.3

Two

1.4

u

2.1

1.7

1.7

1.7

2.3

1.1

2.8

2.0

1.7

Three

1.7

u

2.6

2.9

1.7

u

3.9

u

4.6

u

1.8

One

TABLE 3. Percentage of ever-married women who received three or more antenatal

India	53.5	71.1	52.6	34.3	25.3	1.4	2.1	2.8
Andhra Pradesh	86.0	92.5	86.7	79.3	65.9	1.1	1.2	1.4
Assam	40.2	62.8	41.5	22.1	15.0	1.5	2.8	4.2
Bihar	16.8	37.7	15.7	7.3	6.3	2.4	5.1	6.0
Chhattisgarh	54.5	69.6	55.8	44.6	37.9	1.2	1.6	1.8
Gujarat	68.8	81.0	66.6	47.4	30.8	1.2	1.7	2.6
Haryana	60.7	71.4	54.8	32.9	20.0	1.3	2.2	3.6
Jharkhand	35.5	61.8	39.7	21.5	21.5	1.6	2.9	2.9

Karnataka 79.2 87.6 80.2 62.0 52.7 93.7 94.4 91.0 u u Madhya Pradesh 41.6 58.7 45.8 28.2 22.6

74.2

70.0

64.2

41.0

92.2

24.4

33.6

63.5

Notes: u=unavailable because of small sample size (N<30). All differences in state comparisons across

86.7

79.5

84.2

59.4

99.2

41.5

58.9

81.7

deprivation groups are significant, according to chi-square tests.

76.1

62.6

76.3

41.7

95.8

27.3

46.5

63.0

Kerala

Orissa

**Punjab** 

Rajasthan

Tamil Nadu

Uttar Pradesh

Uttarakhand

West Bengal

Maharashtra

None