

**TABLE 1. Odds ratios from logistic and multinomial logistic analyses examining relationship of selected measures and interactions with pregnancy termination and pregnancy termination method**

Measure	Model 1‡	Model 2§		Model 3‡
	Whether pregnancy terminated	Whether pregnancy terminated by safer method (relative to not terminated)	Whether pregnancy terminated by less-safe method (relative to not terminated)	Whether termination was by safer method (relative to less-safe method)
<b>Maternal age</b>				
≤15	3.20***	3.51***	2.72**	1.34
16–17	1.51***	1.51**	1.52*	0.98
18–19	1.07	1.12	0.99	1.17
20–24 (ref)	1.00	1.00	1.00	1.00
25–29	1.51***	1.69***	1.23*	1.45**
30–34	3.10***	3.58***	2.39**	1.61***
35–39	6.67***	7.71***	5.16**	1.51***
40–44	14.05***	16.41***	10.71**	1.50**
≥45	26.11***	30.54***	20.02**	1.42†
<b>Maternal education</b>				
None (ref)	1.00	1.00	1.00	1.00
1–5 yrs.	1.24***	1.35***	1.07	1.45***
6–10 yrs.	1.37***	1.53***	1.11	1.80***
11–16 yrs.	1.27*	1.43*	0.93	2.17**
Unknown	1.28**	1.34**	1.19	1.14
<b>Time period x area</b>				
1989–1993 x comparison area (ref)	1.00	1.00	1.00	1.00
1994–1998 x comparison area	1.25***	1.63***	0.96	1.68***
1999–2003 x comparison area††	1.79***	2.96***	0.85†	3.56***
2004–2008 x comparison area	1.35***	2.35***	0.52**	4.54***
1989–1993 x MCH-FP area	0.66***	0.67***	0.66***	0.97
1994–1998 x MCH-FP area	0.51***	0.77	0.30**	2.51***
1999–2003 x MCH-FP area††	0.84**	1.37***	0.41**	3.24***
2004–2008 x MCH-FP area	0.84**	1.48***	0.30**	4.84***
Log likelihood	–19,576.9		–22,648.3	–3,056.5

\*p<.05. \*\*p<.01. \*\*\*p<.001. †p<.10. ‡Logistic regression. §Multinomial logistic regression. ††Data not available for 2001. Notes: Models 1 and 2 include all eligible pregnancies; model 3 is restricted to pregnancies that were terminated. ref=reference category. MCH-FP=Maternal and Child Health–Family Planning.